



PESTICIDE INFORMATION SHEET

2025

GROWER ORIGIN	CANADA
CROP	BLUEBERRY
MARKET DESTINATION	CANADA/USA/JAPAN/KOREA/AUSTRALIA
EFFECTIVE DATE	Feb 15 -2025

APPROVED FOR USE			
TRADE NAME	TYPE	PHI	COMMENTS
Actara	Insecticide	3 Days	Post Bloom Only. Crop Renovation Required After Harvest (Canada Only)
Admire	Insecticide	3 Days	Post Bloom Only. Crop Renovation Required After Harvest (Canada Only)
Alias	Insecticide	14 Days	Post Bloom Only. Crop Renovation Required After Harvest (Canada Only)
Altacor Max	Insecticide	1 Day	Do not apply during bloom.
Assail	Insecticide	1 Day	Canadian Growers to Follow 7 Day US PHI. Do not apply during bloom.
Bartlett Supeior	Insecticide	Late Winter	Do not use during bloom.
Bioprotec	Insecticide	0 Days	
Capture	Insecticide	3 Days	Emergency Registration, Expires Sept 30 2022. Do not use during bloom.
Closer	Insecticide	1 day	Do Not Use During Bloom. Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Confirm	Insecticide	14 Days	Use prior to bloom only Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Cormoran	Insecticide	8 Days	Do not use during bloom.
Cyclaniliprole	Insecticide	1 Day	Do not use during bloom.
Danitol	Insecticide	3 Days (15 Days Hand Harvest Canada)	Do not use during bloom.
Decis	Insecticide	Do not use after April 15th	Do not use after april 15th. Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Delegate	Insecticide	3 Days	Do not apply during bloom. 3 Days PHI For Caterpillars
Dipel	Insecticide	0 Days	
Entrust	Insecticide	1 Days	Do not use during bloom. 3 Days PHI For Caterpillars
Exirel	Insecticide	3 Days	Do not apply during bloom.
Foray	Insecticide	0 Days	
Fulfill	Insecticide	85 Days	Use prior to bloom only Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Harvanta	Insecticide	1 Day	Do not use during bloom.
Imidan	Insecticide	15 Days	Do Not Use During Bloom. US Growers follow Canada PHI of 15 days.
Intrepid	Insecticide	7 Days	
Kopa	Insecticide	0 Days	Do not use during bloom.
Lagon	Insecticide	21 Days(Maggot) Post Harvest (SWD)	Toxic to bees. If raspberries are present in field ; contact Berryhill before use. Post harvest only.
Lime Sulphur	Insecticide	Apply During Dormancy	Do Not Apply During Bloom
Malathion	Insecticide	1 Day	Do not use during bloom.
Movento	Insecticide	7 Days	Do Not Use During Bloom. Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Purespray Green Spray	Insecticide	0 Days	Do Not Use During Bloom
Pyganic	Insecticide	0 Days	
Rimon	Insecticide	8 Days	Use prior to bloom only Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Ripcord	Insecticide	2 Days	Do not apply during bloom.
Sivanto	Insecticide	3 Days	Do Not Apply During Bloom
Sluggo	Insecticide	N/A	Slug Bait. Do Not Use During Bloom
Success	Insecticide	1 Day	Do Not Apply During Bloom. 3 Days PHI For Caterpillars
Suffoil-X	Insecticide	12 Hours	Do Not Apply During Bloom
UP-Cyde	Insecticide	2 Days	Do not apply during bloom.
Aim	Herbicide	1 Day	US growers follow CAN PHI of 1 day.
Alion	Herbicide	Fall or early spring only	Fall or early spring only.
Arrow	Herbicide	14 Days	
Authority	Herbicide	Use prior to bloom	Use prior to bloom.
AXXE	Herbicide	0 Days	Bioherbicide.
Beloukha	Herbicide	1 Day	
Callisto	Herbicide	Pre bloom or post harvest use only	Pre bloom or post harvest use only.
Casoron	Herbicide	Late Winter/Fall	Apply in late winter fall.
Centurion	Herbicide	14 Days	
Chateau	Herbicide	7 Days	
Devrinol	Herbicide	70 Days	Spring or fall use.
Dual II Magnum	Herbicide	30 Days	US growers follow CAN PHI of 30 days.
Eragon	Herbicide	Dormancy application only	Apply after leaf drop and before breaking dormancy
Ignite	Herbicide	14 Days	

Lontrel	Herbicide	45 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Matrix	Herbicide	30 Days	Do not apply during bloom.
Princep	Herbicide	Late Fall to Early Spring	Established Plantings Only
Prism	Herbicide	30 Days	Do not apply during bloom. Can growers follow USA PHI of 30 days.
Sandea	Herbicide	14 Days	
Select Max	Herbicide	14 Days	
Serene	Herbicide	0 Days	
Sinbar	Herbicide	80 Days	US growers follow Canada PHI of 80 days.
Velpar	Herbicide	90 Days	Bluecrop only. CAN growers follow US PHI of 90 days.
Aliette WDG/Systemic	Fungicide	1 Day	
Allegro	Fungicide	30 Days	
Bravo / Echo	Fungicide	54 Days	US growers follow Canada PHI of 54 days, Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Bumper/Fitness	Fungicide	60 Days	
Cabrio	Fungicide	1 Day	
Cantus	Fungicide	0 Days	
Captan	Fungicide	2 Days	US growers must follow CAN PHI of 2 days.
Cevya	Fungicide	0 days	
Confine Extra	Fungicide	1 Day	US growers follow Canada PHI of 1 day.
Copper Spray	Fungicide	2 Days	US growers follow Canada PHI of 2 days.
Cueva	Fungicide	1 Day	
Diplomat	Fungicide	0 Days	
Double Nickel	Fungicide	0 Days	
Echo	Fungicide	54 Days	US growers follow Canada PHI of 54 Days, Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Elevate	Fungicide	1 Day	US growers follow Canada PHI.
Fontelis	Fungicide	0 Days	
Indar	Fungicide	30 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Inspire Super	Fungicide	7 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field. Canadian growers follow USA 7 day PHI
Kenja/Isofetamid	Fungicide	7 Days	
Luna Privilege	Fungicide	0 Days	
Luna Tranquility	Fungicide	0 Days	
Maestro	Fungicide	2 Days	
Merivon	Fungicide	See Label	
Metastar 2E	Fungicide	80 Days	US Growers should follow Canadian PHI of 80 days.
Mission	Fungicide	60 Days	
MustGrow	Fungicide	Pre-Plant	
Orondis Gold	Fungicide	80 Day	
OxiDate 2.0	Fungicide	0 Days	
Phostrol	Fungicide	0 Days	
Pivot	Fungicide	60 Days	
Pristine	Fungicide	0 Days	
Problad	Fungicide	0 Days	
Proline	Fungicide	7 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Propulse	Fungicide	7 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Quash	Fungicide	8 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Quilt	Fungicide	60 Days	All Growers Follow 60 day PHI
Regalia (Maxx)	Fungicide	0 Days	
Ridomil Gold 480EC/SL	Fungicide	80 Days	US Growers should follow Canadian PHI of 80 days.
Scala	Fungicide	0 Days	
Sercadis	Fungicide	0 Days	
Serenade Opti/Max	Fungicide	0 Days	
Serifel	Fungicide	0 Days	
Switch	Fungicide	1 Day	US Growers follow Canada PHI of 1 day.
Taegro 2	Fungicide	0 Days	
Tilt	Fungicide	60 Days	US Growers follow Canada PHI of 60 days.
Timorex Gold	Fungicide	2 Days	
Topas	Fungicide	60 Days	US Growers follow Canada PHI of 60 days.
Vegol Crop Oil	Fungicide	0 Days	
Ground Force	Rodenticide	0 Days	Rodent Control
Ramik Green	Rodenticide	0 Days	Rodent Control
Avian Migrate	Bird Control	0 Days	

NOT APPROVED			
2,4-D	Do Not Use	30 Days	Bird Control.
Basagran	Do Not Use	25 Days	
Beleaf	Do Not Use	0 Days	
Capture	Do Not Use	3 days	
Concept	Do Not Use	14 Days Prior to Bloom	
Credit	Do Not Use	30	Use prior to bloom only Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Cygon	Do Not Use	Post Harvest Only	US growers follow Canada PHI of 30 days.
Endivior	Do Not Use	7 Days	
Ferbam	Do Not Use	40 Days	Toxic to bees. Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field . Post harvest only.
Funginex	Do Not Use	60 Days	
Goal	Do Not Use	50 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
Kanemite	Do Not Use	1 day	
Miravis Neo	Do Not Use	30 Days	
Miravis Prime	Do Not Use	1 Day	
Nova	Do Not Use	1 Day	
Poast (Ultra)	Do Not Use	30 Days	
Roundup	Do Not Use	N/A	CAN growers follow US PHI of 30 days.
Sencor	Do Not Use	2 Years	
Sevin	Do Not Use	7 Days	Contact Berryhill prior to use if sprayer is used for raspberries or raspberries are present in field.
StartUp	Do Not Use	30 Days	Do Not Use During Bloom. Canadian growers follow usa PHI of 7 days.
Touchdown	Do Not Use	30 Days	
Vantage	Do Not Use	30 Days	US Growers follow Canada PHI of 30 days.
Venture	Do Not Use	15 Days	
Products Registered In CANADA: Not Likely To Exceed Canada/USA MRL			
Use According To Label Instructions And Follow Any Additional Comments			
No Substitutions			
Disclaimer: The information provided in this list is for general information only, and is not intended as a substitute for the actual product label. The list is believed to be accurate as of the date of this document; however labels are subject to change. With regards to PHI, 1 day is equal to 24 hours and so hours after application and not calendar days is to be considered. Although due care and skill has been exercised in the preparation and compilation of the information and data in this list, Berryhill Foods Inc. does not warrant it's accuracy, completeness, currency or suitability for any purpose. To the maximum extent permitted by law, Berryhill Foods Inc. disclaims all liability, including liability for any negligence, loss, damage, injury or cost of expense incurred by any person as a result of accessing, using or relying upon any of the information or data set out in this document.			